

TO ALL TO WHOM THESE PRESENTS SHALL COME:

Penusylvania Agricultural Experiment Station

MUSICIES, THERE HAS BEEN PRESENTED TO THE

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED DISTINCT VARIETY OF SEXUALLY REPRODUCED, OR TUBER PROPAGATED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED TO BE ENTITLED TO A CERTIFICATE OF PLANT VARIETY PROTECTION UNDER THE LAW.

NOW, THEREFORE, THIS CERTIFICATE OF PLANT VARIETY PROTECTION IS TO GRANT UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLICANT(S) FOR THE TERM OF TWENTY YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EXCLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT, OR IMPORTING IT, OR EXPORTING IT, E CONDITIONING IT FOR PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR CONDITIONING IT PROPAGATION, OR STOCKING IT FOR ANY OF THE ABOVE PURPOSE, OR USING IT IN PRODUCING A HYBRID OR LENT VARIETY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. IN THE STATES SEED OF THIS VARIETY (I) SHALL BE SOLD BY VARIETY NAME ONLY AS A CLASS OF CERTIFIED SEALL CONFORM TO THE NUMBER OF GENERATIONS SPECIFIED BY THE OWNER OF THE RIGHTS. (84 STAT. 1542, 200).

BENTGRASS, CREEPING

'Penn A-1'

In Testimon Mucros, I have hereunto set my hand and caused the seal of the Mant Mariety Arstection Office to be affixed at the City of Washington, D.C. this eighth day of October, in the year two thousand and four.

Attest:

Commissioner
Plant Variety Protection Office

Agriculture

Pennsylvania Agricultural Experiment Station

U.S. DEPARTMENT OF AGRICULTURE

AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE The following statements are made in accordance with the Privacy Act of 1974 (5 U.S.C. 552a) and the Paperwork Reduction Act (PRA) of 1995. Application is required in order to determine if a plant variety protection

certificate is to be issued (7 U.S.C. 2421). Information is held confidential

			-
PPLICATION FOR PLANT	VARIETY	PROTECTION	CERTIFICATE

(Instructions and information collection burden statement on reverse) 1. NAME OF APPLICANT(S) (as it is to appear on the Certificate)

2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER

until certificate is issued (7 U.S.C. 2426).

Penn A-1

3. VARIETY NAME

4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country)

0217 Agricultural Administration Building The Pennsylvania State University University Park, PA 16802

5. TELEPHONE (include area code)

814-865-5410

PVPO NUMBER 9700046

Greeping Rentgrass

FOR OFFICIAL USE ONLY

6. FAX (include area code)

814-863-7905

November 26, 199

DATE

7. GENUS AND SPECIES NAME

Agrostis palustris

8. FAMILY NAME (Botanical)

Gramineae

FILING AND EXAMINATION FEE: • 2450 ºº

8. CROP KIND NAME (Common name)

Creeping Bentgrass

10. IF THE APPLICANT NAMED IS NOT A "PERSON", GIVE FORM OF ORGANIZATION (corporation, partnership, association, etc.) (Common name) Land Grant University

11. IF INCORPORATED, GIVE STATE OF INCORPORATION

12. DATE OF INCORPORATION

DATE November 26, 199

13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE IN THIS APPLICATION AND RECEIVE ALL PAPERS

Dr. Charles R. Krueger Bruce A. Mc Pheron, Ph.D. Associate Dean

0217 Agricultural Administration Building University Park, PA 16802

15. FAX (include area code)

14. TELEPHONE linclude area code!

16. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow instructions on reverse)

- a. 🔀 Exhibit A. Origin and Breeding History of the Variety
- b. 🔀 Exhibit B. Statement of Distinctness
- c. 🛛 Exhibit C. Objective Description of the Variety
- d. 🛛 Exhibit D. Additional Description of the Variety (Optional)
- e. X Exhibit E. Statement of the Basis of the Applicant's Ownership
- f. 🔯 Voucher Sample (2,500 viable untreated seeds or, for tuber propagated varieties varification that tissue culture will be deposited and maintained in an approved public repository)
- g. 🏻 Filing and Examination Fee (\$2,450), made payable to "Treasurer of the United States" (Mail to PVPO)

17. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VARIETY NAME ONLY, AS A CLASS OF CERTIFIED SEED? (See Section 83(a) of the Plant Variety Protection Act)

YES (If "yes," answer items 18 and 19 below)

NO (If "no," go to item 20)

19. IF "YES" TO ITEM 18, WHICH CLASSES OF PRODUCTION BEYOND BREEDER SEED?

18. DOES THE APPLICANT SPECIFY THAT SEED OF THIS VARIETY BE LIMITED AS TO NUMBER OF **GENERATIONS?**

FOUNDATION REGISTERED E CERTIFIED

20. HAS THE VARIETY OR A HYBRID PRODUCED FROM THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MARKETED IN THE U.S. OR OTHER COUNTRIES? X YES (If "yes," give names of countries and dates)

USA, November 30, 1995

Commercial Sale

21. The applicant(s) declare that a viable sample of basic seed of the variety will be furnished with application and will be replenished upon request in accordance with such regulations as may be applicable, or for a tuber propagated variety a tissue culture will be deposited in a public repository and maintained for the duration of the certificate.

The undersigned applicant(s) is(are) the owner(s) of this sexually reproduced or tuber propagated plant variety, and believe(s) that the variety is new, distinct, uniform, and stable as required in Section 42, and is entitled to protection under the provisions of Section 42 of the Plant Variety Protection Act.

Applicant(s) is(are) informed that false representation herein can jeopardize protection and result in penalties.

SIGNATURE OF APPLICANT (Owner(s))

X YES

SIGNATURE OF APPLICANT (Owner(s))

Associate Dean

CAPACITY OR TITLE

Charles R. Krueger

10/17/96

CAPACITY OR TITLE

NAME (Please print or type)

DATE

STD-470 (03-96) (Previous editions are to be destroyed)

INSTRUCTIONS

GENERAL: To be effectively filed with the Plant Variety Protection Office (PVPO), ALL of the following items must be received in t PVPO: (1) Completed application form signed by the owner; (2) completed Exhibits A, B, C, E; (3) at least 2,500 viable untreat seeds, or for tuber reproduced varieties verification that a viable (in the sense that it will reproduce an entire plant) tissue culture we be deposited and maintained in an approved public repository; (4) check drawn on a U.S. bank for \$2,450 (\$300 filing fee at \$2,150 examination fee), payable to "Treasurer of the United States" (See Section 97.6 of the Regulations and Rules of Practice Partial applications will be held in the PVPO for not more than 90 days, then returned to the applicant as unfilled. Mail application at other requirements to Plant Variety Protection Office, AMS, USDA, Room 500, NAL Building, 10301 Baltimore Blvd., Beltsville, M 20705-2351. Retain one copy for your files. All items on the face of the application are self explanatory unless noted below Corrections on the application form and exhibits must be initialed and dated. DO NOT use masking materials to make corrections. a certificate is allowed, you will be requested to send a check payable to "Treasurer of the United States" in the amount of \$300 files issuance of the Certificate.

Plant Variety Protection Office Telephone: (301) 504-5518

ITEM

- 16a. Give: (1) the genealogy, including public and commercial varieties, lines, or clones used, and the breeding method;
 - (2) the details of subsequent stages of selection and multiplication;
 - (3) evidence of uniformity and stability; and
 - (4) the type and frequency of variants during reproduction and multiplication and state how these variants may be identified.
- 16b. Give a summary of the variety's distinctness. Clearly state how this application variety may be distinguished from all othe varieties in the same crop. If the new variety is most similar to one variety or a group of related varieties:
 - (1) identify these varieties and state all differences objectively;
 - (2) attach statistical data for characters expressed numerically and demonstrate that these are clear differences;
 - (3) submit, if helpful, seed and plant specimens or photographs (prints) of seed and plant comparisons which clearly indicate distinctness.
- 16c. Exhibit C forms are available from the PVPO for most crops; specify crop kind. Fill in Exhibit C (Objective Description of Variety) form as completely as possible to describe your variety.
- Optional additional characteristics and/or photographs. Describe any additional characteristics that cannot be accurately conveyed in Exhibit C. Use comparative varieties as is necessary to reveal more accurately the characteristics that are difficult to describe, such as plant habit, plant color, disease resistance, etc.
- 16e. Section 52(5) of the Act requires applicants to furnish a statement of the basis of the applicant's ownership. An Exhibit E form is available from the PVPO.
- 17. If "Yes" is specified (seed of this variety be sold by variety name only, as a class of certified seed), the applicant may NOT reverse this affirmative decision after the variety has been sold and so labelled, the decision published, or the certificate issued. However, if "No" has been specified, the applicant may change the choice. (See Regulations and Rules of Practice, Section 97.103).
- 20. See Sections 41, 42, and 43 of the Act and Section 97.5 of the regulations for eligibility requirements.

NOTES: It is the responsibility of the applicant/owner to keep the PVPO informed of any changes of address or change of ownership or assignment during the life of the application/certificate. There is no charge for filing a change of address. The fee for filing change of ownership or assignment is specified in Section 97.175 of the regulations. (See Section 101 of the Act, and Section 97.130, 97.131, 97.175(h) of Regulations and Rules of Practice.)

To avoid conflict with other variety names in use, the applicant should check the variety names proposed by contacting: Seed Branch, AMS, USDA, Room 213, Building 306, Beltsville Agricultural Research Center-East, Beltsville, MD 20705.
Telephone: (301) 504-8089.

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, Jamie L. Whitten Building, Washington, D.C. 20250. When replying, refer to OMB No. 0581-0055 and form number in your letter. Under the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

The U.S. Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (2021 720-2791. To file a complaint, write the Socretary of Agriculture, U.S. Department of Agriculture, Washington, D.C. 20250, or call (2021 720-7327 (voice) or (2021 720-1127 (TDD). USDA is an equal employment opportunity employer.

Attachment for Application Item 20.

	PSU Release ¹	PSU PVP Application ²	PVP Application Date ³	First Commercial Sale
Penn A-1	08-29-95	10-17-96	11-26-96	11-30-95
Penn A-2	08-29-95	10-17-96	12-19-96	02-27-96
Penn A-4	08-29-95	10-17-96	12-19-96	01-24-96
Penn G-1	08-29-95	02-21-97	02-26-97	05-20-96
Penn G-2	08-29-95	10-17-96	12-12-96	01-10-96
Penn G-6	08-29-95	10-17-96	01-02-97	01-24-96
Seaside II	08-29-95	10-17-96	12-12-96	01-26-96

¹PSU Release - Date Pennsylvania Experiment Station Seed Committee approved variety release

²PSU PVP Application - Date on PSU PVP application form

³PVP Application - Application date of variety by PVP office as listed in PVP Official Journal, Vol. 25, December 1997

Penn A-1

Origin and Breeding History of Penn A-1, A-2, A-4, G-1, G-2, G-6

The objective of this varietal breeding program was to develop creeping bentgrasses exhibiting superior putting green turf over existing varieties. Original parental material was selected from segregated patches of bent, 12 to 18 inches in diameter, on some greens at the Augusta National Golf Club, Augusta, Georgia in the spring of 1984. These segregates attracted attention because they were vigorous and dense, very fine textured, and had a more upright growth habit of individual plant tillers. The growth habit was unique because the selected segregates did not spike (raise up) from golfer's metal-spiked golf shoes. Their ability to spread at a closer than normal height of cut used for fast greens was also indicative of potential heat tolerance in a hot, humid golf course environment. The value of non-spiking features was proven in subsequent years by the banning of golf shoes with metal spikes on most American golf courses, except for professional tournaments.

There were two groups of selections by the breeder, the 'G' and 'A' series. There were eight 'G' selections (G-1 to G-8) from two greens on the Par 3 course originally seeded to Penneagle creeping bent, and six selections (A-1 to A-6) from four greens on the main course originally seeded to Penncross creeping bent.

The breeding method used was a polycross procedure. The single parent selection of Penn A-1 was crossed with experimental selections of Penn G followed by two generations of crossing selected sib plants of Penn A-1.

Both 'G' and 'A' series selections were cloned into eight plants, pot planted, and induced to flower in growth chambers for six weeks. Isolated crossing blocks were established in the greenhouse for each 'G' and 'A' line in December 1984. Due to near self-sterility of bent, 'A' line plants were used as male crossing parents for 'G' lines, and conversely, 'G' plants used as male parents for the 'A' lines. Seed was produced from all crosses in March. Next 250 plants from bulk seed of each 14 lines (G-1, G-2, G-3, G-4, G-5, G-6, G-7, G-8 and A-1, A-2, A-3, A-4, A-5, and A-6) were nursery space planted in isolated field blocks in August 1985.

The first cycle of reselection began in the spring of 1986. First plants to be chemically rogued were those lacking in vigor coming out of winter. Majority of plants had a more upright growth with short stolons in keeping with their putting green selections. There were a few semi-decumbent types with longer stolons and dense ball types. The emphasis was on selecting the most uniform in upright growth habit, vigor, and uniformity in pre-anthesis flowering. On this basis, 30 to 50 plants were selected from each block of A-1, A-2, A-4, G-1, G-2, and G-6. All other 'A' and 'G' lines were discarded due to a combination of segregation and lack of vigor uniformity.

Seed was harvested from selected lines (G-1, G-2, G-6, A-1, A-2, and A-4) and used for small turf plots which confirmed the fine, dense qualities of original parents. The

N-9700046

selected clones of the first cycle were pot planted, cloned, and again induced to flower in growth chambers to save a year. Plants of each line were in isolated greenhouse crossing blocks. Crossing in this cycle was confined to the siblings of each line with no other pollen source. From the crossed plants of each line, 300 seedlings were grown for field planting for the next cycle of reselection in 1987.

The second cycle of reselection in 1987 consisted of arbitrarily selecting 40 to 50 plants as a reasonable number to work with and with the main criterion of selecting an upright growth habit eliminating all but very short stolon types, uniformity of size and vigor, and flowering. Confining the pollen source in this crossing within the A-1 parents resulted in increasing the population of desired growth habit types and reducing the longer stolon types to a few. Most of the "off-types" were limited to the dense ball types with few flowering heads.

Forty clones of each G and A were recloned to three plants and sent to Oregon for further evaluation and reselection under production state conditions where growth is more robust than in Pennsylvania and where commercial production would eventually take place. Following observations, 20 clones each were selected as potential Breeders parents. Seed of each G and A line was used to plant half acre observation, seed yields, and management trials. Satisfactory uniformity and stability of the G and A bent varieties from Oregon grown first generation seed was ascertained by inspection in both the vegetative and flowering stages of growth by the breeder, pure seed testing personnel, and Oregon certification. Variants consisted of a few more spreading decumbent types and the non-spreading ball types. The decumbent type seed heads protrude laterally at plant perimeters and the ball types produce few or no flowers. The decumbent variants for A-1 were calculated to constitute 0.17% based on 49 plants per half acre of an estimated population of 39,000 plants.

By agreement with the Bentgrass Growers Association with proprietary rights, all observed variants shall be chemically rogued to maintain varietal purity and stability in order to maintain a certified stand life of five seed yield years. To further maintain varietal purity, the commercial production of Penn G and A bents shall be limited to a two generation system, Breeder and Certified; the only varieties limited to only two generations.

Three cycles of five year stands of commercial plantings have shown that Penn A-1 is a uniform and stable variety to the satisfaction of the breeder and Oregon Certification with no further reselection deemed necessary.

Breeder seeds of Penn A-1 has been maintained and produced by Pure Seed Testing in Hubbard Oregon since 1994. Approval of the variety name has been cleared by the Seed Branch on May 21, 2003.

Variety Distinction of Penn A-1

A PVP nursery was established at University Park, Pennsylvania in 1994 consisting of 21 creeping bent varieties with three replications of 25 spaced plants. Included were Penn A-1, plus six new Penn State varietal releases Penn G-1, G-2, G-6, A-2, A-4, Seaside II, and 13 commercial varieties. Data were collected in 1995 as shown in Table 1. This nursery was discarded due to loss of land.

A second PVP nursery was established at the Pure Seed Testing Research Farm near Hubbard, Oregon in 1995. The purpose was to evaluate plants in the location of major bentgrass commercial production where growth greatly exceeds the environment in the Northeast. It consisted of the above experimentals and 12 commercial varieties with four replications of 25 spaced plants. This test was maintained and data collected and analyzed by Pure Seed personnel after the original application for PVP. These data are shown in Tables 2 and 3. Varieties significantly different by years from Penn A-1 are summarized in Table 4. In this form, the most obvious differences and similarities are easily discerned.

Penn A-1 may be most easily distinguished from other bentgrass varieties tested predominately by plant height and panicle length based on morphological characters measured in three years of testing. It is most similar to Penn G-2. Another difference was the following:

Characteristic	Penn A-1
Ligule shape at apex	80% Acute
	20% Truncate

<u>vs</u>	Penn G-2
	20% Acute
-	80% Truncate

U.S. Department of Agriculture Agricultural Marketing Service Science Division Beltsville, Maryland 20705

OBJECTIVE DESCRIPTION OF VARIETY BENTGRASS (Agrostis spp.)

Name of Applicant(s) Pennsylvania Agricultural Experiment Station	Variety Name or Temporary Designation
Address (Street and No. or R.F.D. No, City, & ZIP Code)	Penn A-1 (A-1)
0217 Agricultural Administration Bldg.	FOR OFFICIAL USE ONLY
The Pennsylvania State University University Park, PA 16802	PVPO Number 7 0 0 0 4 6
Place numbers in the boxes (e.g. 089) for the typical plants of this variety. The symb	e characters that best describe ool & indicates decimal.
COMPARISON VARIETIES FOR USE BELOW	
l= Astoria 2= Exeter 3= Highland 4= Seaside 7= Astra 8= Other Lopez	5= Penncross 6= Kingstown 9-Pennlinks 10-Southshore
	ping A. stolonifera (A.palustris m bent A. canina ssp. montana
5- Red top A. gigantea	
5- Red top A. gigantea 2. ADAPTATION: (0- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest	2= Adapted)
5- Red top A. gigantea 2. ADAPTATION: (0- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central	2- Adapted) 2 Pacific N. W.
5- Red top A. gigantea 2. ADAPTATION: (0- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest	2- Adapted) 2 Pacific N. W. arieties
5- Red top A. gigantea 2. ADAPTATION: (O= Not Tested, l= Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest 3. MATURITY (At first anthesis): Use comparison v	2- Adapted) 2 Pacific N. W. arieties ,06 Days later than 8
5- Red top A. gigantea 2. ADAPTATION: (O- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest 3. MATURITY (At first anthesis): Use comparison v 0. 7 Days earlier than 5 , Maturity same as 1. 4. HEIGHT (Average of longest 10 shoots from soil	2- Adapted) 2 Pacific N. W. arieties ,06 Days later than 8 surface to top of head):
5- Red top A. gigantea 2. ADAPTATION: (O- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest 3. MATURITY (At first anthesis): Use comparison v 0. 7 Days earlier than 5 , Maturity same as 1 4. HEIGHT (Average of longest 10 shoots from soil 4. 8 Cm Height (at maturity)	2- Adapted) 2- Adapted) 2- Pacific N. W. arieties ,06 Days later than 8 surface to top of head): ter than 4 comparison same as 10
5- Red top A. gigantea 2. ADAPTATION: (O- Not Tested, 1- Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest 3. MATURITY (At first anthesis): Use comparison v 0. 7 Days earlier than 5 , Maturity same as 1 4. HEIGHT (Average of longest 10 shoots from soil 4. 8 Cm Height (at maturity)	2- Adapted) 2 Pacific N. W. arieties ,06 Days later than 8 surface to top of head): ter than 4 comparison variety
5- Red top A. gigantea 2. ADAPTATION: (O= Not Tested, l= Not Adapted, 2. Northeast 2 Southeast 2 North Central 2. Other (Specify) Southwest 3. MATURITY (At first anthesis): Use comparison v 0. 7 Days earlier than 5 , Maturity same as 1 4. HEIGHT (Average of longest 10 shoots from soil 4. Meight (at maturity) O. 7 Cm Shore	2- Adapted) 2- Adapted) 2- Pacific N. W. arieties ,06 Days later than 8 surface to top of head): ter than 4 comparison variety

Bentgrass - 2 -

6. VEGETATIVE REPRODUCTION:	
Rhizomes 1- Absent 2- Present Rhizomes 1- Absent 2- Present Rhizomes 1- Absent 2- Present	2 Stolons 1= Absent 2= Present
7. LEAF BLADE:	
3 Color: l= Yellowish Green (Cohansey) 3= Green (Exeter) 5= Bluish Green (Highland) 1 Texture: (fineness)	2- Light Green (Washington) L= Dark Green (Kingstown, Tracenta 6- Other (Specify)
l= Very fine (Kingstown) 3= Medium fine (Astoria) 5= Medium coarse (Virginia)	2= Fine (Exeter) 4= Medium (Seaside) 6= Coarse (Vermont)
Stomatal density upper leaf surface (Nu	mber/mm ²)
The same of the sa	∫% Rough
	% Rough
	% Rough
Mm Width (Average of 10)	arrower than []
Widt	h same as
	ider than
3 3 M W 11 41 (m	ength (Flag leaves)
8. LEAF SHEATH:	
Anthocyanin: 1= Absent 2= Present	% Red sheaths
9. LIGULE (Lower and middle leaves):	
Shape at Apex: [8]0] & Acute [] & Ro	ounded 20 4 r
5 Other (Specify)	A Truncate
Pubescence: T1010 % Glabrous	
Margins: 130 % Entire 770 %	Fronthad
7 Other (Specify)	Toothed
Mm Length	

10. LEMMA:
Shape: Sh
411 Mm Width 1A7 Mm Length (exclusive of awn)
Color: The Buff 1010 % Silvery The Coher (Specify)
Surface 100 % Glossy Dull
Texture: 100 % Smooth % Punctate
Pubescence: 100 % Glabrous % Sparse Copious
Basal Hairs: 100 % Absent
Short III % Long
Appressed Ascending Spreading
Awns: 100 % Absent Few Many
Awn-pointed # Short # Long
Straight Geniculate
Awn Insertion on Lemma:
TA DANTOTO
11. PANICLE:
Type (in anthesis): 5 0 % Open 5 5 0 % Compact
Anthocyanin: 100 % Absent 7 % Present
Branches in Anthesis: 8.0 % Appressed 210 % Ascending % Spreading
Branches in Fruit: 8 0 % Appressed 20 % Ascending % Spreading
Branch Surface: 19 0 % Smooth 11 0 % Scabrous
12. SEED:
47 1 3rams per 1000 seed
13. SPRING GREEN UP:
2 l= Early (Exeter) 2= Medium (Astoria) 3= Late (Kingstown)

Bentgrass - 4 -

2 Cold 2 Heat	Susceptible 2= Resistant)	tested, 1	: (O= Not	l resistanci	14. ENVIRONMENTA
Red Leaf Spot - Drechslera erythrospila Helminthosporium Leaf Spot Bipolaris scrokiniana				· ·	
Red Leaf Spot - Drechslera erythrospila Helminthosporium Leaf Spot (Bipolaris scrokiniana) Melting Out - Drechslera poae (Helminthosporium vagans) Pythium Blight - (P. sphanidermatum) Pythium Blight (P. ultimum) Pusarium Blight (F. roseum) Pythium Blight (P. ultimum) Pusarium Blight (F. roseum) Pythium Blight (E. tricinctum (P. nivale) Ophiobolus Patch (Pink Snow Mold) Powdery Mildew (Erysiphe gramum (P. nivale) Ophiobolus Patch (O. graminis) Stripe Smut (Ustilago stripe Smut (Ustilago stripe Stripe Smut (Ustilago stripe Stripe Smut (P. coronata) Red Thread (Corticium fuciforme) Typhula Blight (Snow Scald) (T. incarnata) Red Thread (Corticium fuciforme) Other Stem Rust (Puccinia graminis) Other Leaf Rust (P. poae-nemoralis) Other Crown Rust (P. coronata) Leaf Rust (P. poae-nemoralis) Other Garden Chafer	ptible 2= Resistant):	l= Susce	Not teste	STANCE (0=	15. DISEASE RESI
Melting Out - Drechslera poae (Helminthosporium vagans) I Pythium Blight - (P. aphanidermatum) Pythium Blight (F. roseum) Pythium Blight (P. ultimum) Pother (F. roseum) Existance (P. unitimum) Pythium Blight (P. ultimum) Pythium Blight (P. ultimum) Pythium Blight (P. ultimum) Pother (F. roseum) Existance (P. unitimum) Pythium Blight (P. ultimum)					
(Helminthosporium vagans) [Pythium Blight - (P. sphanidermatum)	(Pinologia	ospita []			
(Helminthosporium vagans) I Pythium Blight - (P. aphanidermatum) Pythium Blight (F. roseum) Fusarium Blight (F. roseum) Fusarium Blight (F. roseum) Fusarium Blight (F. roseum) Fusarium Blight (F. tricinctum) Fusarium Patheta	Coloratia sorokiniana)	9	a poae	- Drechsler	Melting Out
Fusarium Blight (F. roseum) Pusarium Blight (F. roseum) Pusarium Patch (Pink Snow Mold) (F. nivale) Ophiobolus Patch (O. graminis) Gopper Spot (Gloeocercospora sorghi) Red Thread (Corticium fuciforme) Stem Rust (Puccinia graminis) Leaf Rust (P. poae-nemoralis) Crown Rust (P. coronata) Leaf Rust (P. poae-nemoralis) Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) Typhula Blight (Snow Scald) (T. incarnata) Brown Patch (Rhizoctonia solder (P. coronata) Crown Rust (P. coronata) Chher Garden Chafer (Amphimallon solstitialis) Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) Tother Other Crambus Spp.) Tother Tother Collowing characteristics indicate degree of resemblance (D.R.) with one of sollowing numbers; 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character Simila	Solida opot =(Scierotinia homo	agans)	osporium	(Helminth	
Fusarium Blight (F. roseum) [Pusarium Patch (Pink Snow Mold) [F. nivale) [Ophiobolus Patch (O. graminis) [Copper Spot (Gloeocercospora sorghi) [Red Thread (Corticium fuciforme) [Stem Rust (Puccinia graminis) [Leaf Rust (P. poae-nemoralis) [Comper Chafer [Amphimallon solstitialis) [Chinch Bug (Blissus insularis) [Armyworm (Cutworm) [Pseudaletia unipuncta) [Other [Oth	ythium Blight (P. ultimum)	<u>m</u>)	naniderma	ght - (P. ap	Pythium Blig
Powdery Mildew (Erysiphe gra (F. nivale) Powdery Mildew (Erysiphe gra (F. nivale) Stripe Smut (Ustilago strii			eum)	ght (F. rose	Fusarium Bli
Ophiobolus Patch (O. graminis)	•		w Mold)	ch (Pink Sno	2 Fusarium Pat
Copper Spot (Gloeocercospora sorghi) Typhula Blight (Snow Scald) (T. incarnata) (T.			minis)	<u>e)</u> atch (0. gra	Ophiobolus P
Red Thread (Corticium fuciforme) Stem Rust (Puccinia graminis) Leaf Rust (P. poae-memoralis) Crown Rust (P. coronata) Leaf Rust (P. poae-memoralis) Chher Coronata Coronata Other Coronata Coronata		. —			
Stem Rust (Puccinia graminis) Leaf Rust (P. poae-nemoralis) Crown Rust (P. coronata) Leaf Rust (P. poae-nemoralis) Chiner Carden Chafer (Amphimallon solstitialis) Chineh Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) Cother Cother Carden Chafer (Phyllopertha horticola) Webworm (Crambus spp.) Armyworm (Cutworm) (Pseudaletia unipuncta) Cother Cother Carden Chafer (Phyllopertha horticola) Cother (Pseudaletia unipuncta) Cother Cother Cother Carden Chafer (Phyllopertha horticola) Cother (Pseudaletia unipuncta) Cother Cother Carden Chafer (Phyllopertha horticola) Cother (Pseudaletia unipuncta) Cother (Pseudaletia unipuncta) Cother (Pascularity of the summitted variety: For the solid paracteristics indicate degree of resemblance (D.R.) with one of similar variety, 2= Same as, 3= More than, darker or superior, etc. Character Similar Variety D.R. Character Similar Variety D.R. Character Similar Variety D.R. Character Similar Variety D.R. Character Foundh Habit Lopez 3 Leaf Color Pennlinks 2 Pennlinks 2 Heat Resistance Lopez 3 Lopez 3 Lopez 3 Lopez 4 Pennlinks 2 Heat Resistance Lopez	ypnula blight (Snow Scald)	±/			
Leaf Rust (Puccinia graminis) Leaf Rust (P. poae-nemoralis) Crown Rust (P. coronata) Other 16. INSECT RESISTANCE (P. Not tested, 1= Susceptible, 2= Resistant): European Chafer (Amphimallon solstitialis) Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) Other (Pseudaletia unipuncta) 7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character Similar Variety D.R. Ch	rown Patch (Rhizoctonia solar		uciforme)	Corticium f	Red Thread (
Leaf Rust (P. poae-nemoralis) Other 16. INSECT RESISTANCE (D- Not tested, 1= Susceptible, 2= Resistant): European Chafer (Amphinalion solstitialis) (Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) Other (Pseudaletia unipuncta) Other (Pseudaletia unipuncta) Other (Other (O	•		inis)	uccinia gram	Stem Rust (P
Carden Chafer Garden Chafer Garden Chafer Phyllopertha horticola					_
European Chafer (Amphimallon solstitialis) (Chinch Bug (Blissus insularis) (Chinch Bug (Blissus insularis) (Phyllopertha horticola) (Phyllopertha	ther	ه لــا د	alis)	. poae-nemor	Li Lear Must (P.
European Chafer (Amphimallon solstitialis) (Chinch Bug (Blissus insularis) (Chinch Bug (Blissus insularis) (Phyllopertha horticola) (Phyllopertha	le 2m Registent)	Suscentii	tested.	INCE (D- Not	16. INSECT RESISTA
(Amphimallon solstitialis) (Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) (Phyllopertha horticola) Webworm (Crambus spp.) I call the substitute of the	20, 2- Resistant);				P 7
Chinch Bug (Blissus insularis) Armyworm (Cutworm) (Pseudaletia unipuncta) 7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of following numbers; 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character From the following numbers; 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character From Length Similar Variety D.R. Character Fourth Habit Lopez Turf Fineness Lopez Heat Resistance Pennlinks Pennlinks Pennlinks Pennlinks Pennlinks Pennlinks Pennlinks Pennlinks Pennlinks	rden Chafer	L G		er	Liropean Chaf
Armyworm (Cutworm) (Pseudaletia unipuncta) 7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character Similar Variety D. with the condition of the cond			<u>is</u>)	n solstitia	(Amphimallo
Armyworm (Cutworm) (Pseudaletia unipuncta) 7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter Similar Variety D.R. Character Similar Variety D. A. Character	bworm (Crambus spp.)	W	aris)	lissus insu	Conner and (R
(Pseudaletia unipuncta) 7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter	— — — .	_		worm)	Armyworm (Cut
7. GIVE VARIETY(S) THAT MOST CLOSELY RESEMBLE THE SUBMITTED VARIETY: For the following characteristics indicate degree of resemblance (D.R.) with one of following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter	her)		
following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter			•		
following numbers: 1= Submitted variety is less than, lighter, or inferior similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter	CIRMINAL MARTINE	SEMBLE THE	CLOSETY R) THAT MOST	7. GIVE VARIETY(S
similar variety, 2= Same as, 3= More than, darker or superior, etc. haracter					
haracter Similar Variety D.R. Character Similar Variety D. rowth Habit Lopez 3 Leaf Color Pennlinks 2 main Length Panicle Type Pennlinks 2 cold Resistance Pennlinks 2 Heat Resistance Lopez 3					
haracter Similar Variety D.R. Character Similar Variety D. rowth Habit Lopez 3 Leaf Color Pennlinks 2 m Length Panicle Type Pennlinks 2 cld Resistance Pennlinks 2 Heat Resistance Lopez 3	arker or superior etc	re than, d	as, 3= M	7, 2= Same	similar variety
rowth Habit Lopez 3 Leaf Color Pennlinks 2 rowth Habit Lopez 3 Lopez 3 Leaf Color Pennlinks 3 Lopez 3 Leaf Color Pennlinks 4 Leaf Colo					haracter
wm Length Panicle Type Pennlinks Turf Fineness Lopez Pennlinks Heat Resistance Lopez				Lopez	
eed Weight Pennlinks 2 Turf Fineness Lopez 3 Fought Resistance Pennlinks 2 Heat Resistance Lopez 3					
old Resistance Pennlinks 2 Heat Resistance Lopez				·	
rought Registered Lopez		TUTI F		Pennlinka	
I Shoda Basishanad				/	ought Resistance
rown Patch		Moss Re	2	Lopez	
	TEIMILIIKS 3				

Table 1. Morphological Character Measurements¹ 1995

¹Penn State University Breeding Nursery, University Park, PA. Three replications of 25 space plants each.

Table 2. Morphological Character Measurements¹ 1996

							Action 1	***************************************	#
	ì		:		Flag		Flag		Bottom
	Plant		Panicle		Leaf		Leaf		Whorl
	Height		Length		Length		Width		Branches
	(cm)		(cm)	:	(cm)		(mm)		
Seaside II	63.8	Penn A-1	13.1	Penn G-1	9.25	Providence	4.80	Penn A-1	7.3
Seaside	63.1	Seaside	13.0	Seaside II	9.03	Seaside II	4.57	Penn A-2	7.0
Penn A-1	62.5	Southshore	12.8	Seaside	8.90	Lopez	4.53	Procup	7.0
Southshore	60.2	Crenshaw	12.3	Regent	8.47	Crenshaw	4.52	Crenshaw	6.7
Penn G-2	57.1	Cato	12.2	Procup	8.26	Penn A-1	4.45	Lopez	9.9
Crenshaw	54.1	Penn G-1	11.7	Penn A-1	7.98	Cato	4.40	Putter	6.4
Providence	53.1	Penn G-6	11.7	Putter	7.91	_ Procup	4.37	Penneagle	6.4
Penn G-1	53.0	Seaside II	11.6	Penn G-6	7.82	Penncross	4.32	Pennlinks	6.4
Penn A-2	52.2	Penn A-4	11.6	Cato	7.78	Penn G-6	4.17	Penn G-6	6.3
Regent	51.4	Penn G-2	11.4	Penn A-2	7.73	Southshore	4.17	Penncross	6.3
Putter	49.9	Putter	11.2	Crenshaw	7.48	Penn G-2	3.95	Providence	6.3
Lopez	47.5	Pennlinks	11.0	Penn G-2	7.43	Pennlinks	3.83	SR 1020	6.0
Penn A-4	46.7	Regent	11.0	Lopez	7.13	SR 1020	3.82	Cato	5.8
Penneagle	46.0	Penneagle	11.0	Penncross	7.06	Penn A-4	3.77	Penn G-2	5.6
SR 1020	44.3	Procup	10.8	Southshore	7.06	Putter	3.75	Regent	5.6
Procup	44.3	Lopez	10.5	Penneagle	6.87	Penneagle	3.62	Penn G-1	5.6
Penn G-6	43.2	Providence	10.2	SR 1020	98.9	Seaside	3.43	Southshore	5.5
Penncross	43.2	SR 1020	10.0	Providence	89.9	Penn G-1	2.83	Seaside II	5.4
Cato	42.3	Penncross	10.0	Pennlinks	6.50	Penn A-2	2.58	Seaside	4.8
Pennlinks	37.8	Penn A-2	6.6	Penn A-4	5.90	Regent	2.38	Penn A-4	4.6
LSD (0.05)	5.2		0.67		1.56		96.0		0.93

¹Pure Seed Testing Research Farm, Hubbard, Oregon. Four replications of 25 space plants each.

Table 3. Morphological Character Measurements¹ 1997

#	Bottom	Whorl	Branches		5.18	4.95	4.95	4.63	4.62	4.53						4.28	4.17	4.17	4.17	4.00	3.82	3.60	3.45	3.43	0.59
					Penncross	Penneagle	Putter	Penn G-1	Seaside II	Lopez	Penn A-2	Providence	Southshore	Crenshaw	Pennlinks	SR 1020	Regent	Seaside	Penn G-2	Cato	Penn A-4	Penn A-1	Penn G-6	Procup	
	Flag	Leaf	Width	(mm)	4.50	4.27	3.35	3.30	3.23	3.12	3.00	2.95	2.93	2.82	2.82	2.73	2.70	2.70	2.67	2.58	2.58	2.45	2.40	2.37	0.30
100					Seaside	Penneagle	Procup	Penncross	Lopez	Penn G-1	Penn A-1	Putter	Penn A-4	Pennlinks	Penn A-2	Penn G-2	Penn G-6	Cato	Providence	Regent	Crenshaw	Seaside II	SR 1020	Southshore	
ļ	Flag	Leaf	Length	(cm)	7.58	7.05	7.00	86.9	6.97	6.72	89.9	6.55	6.40	6.04	5.77	5.71	5.26	4.87	4.82	4.55	4.35	4.29	3.97	3.83	0.54
					Seaside	Providence	Southshore	Penn G-2	Seaside II	Pennlinks	Crenshaw	Cato	SR 1020	Penn G-1	Penneagle	Penncross	Penn A-1	Procup	Penn A-4	Penn A-2	Lopez	Penn G-6	Regent	Putter	
	:	Panicle	Length	(cm)	12.2	11.3	11.1	10.7	10.4	10.3	10.1	10.0	10.0	10.0	6.6	8.6	9.6	9.4	9.3	9.1	9.1	8.9	8.1	7.9	0.75
				:	Seaside	Pennlinks	Penneagle	Crenshaw	Providence	Seaside II	Southshore	SR 1020	Regent	Cato	Penncross	Penn G-1	Penn G-2	Penn A-2	Penn A-1	Putter	Lopez	Penn A-4	Procup	Penn G-6	
	2	Flant	Height	(cm)	63.2	59.4	55.9	54.6	53.9	53.6	53.5	53.4	53.0	52.7	52.2	51.6	50.1	49.8	49.4	49.1	47.6	47.4	46.8	46.3	2.9
				:	Seaside	Penneagle	Penn G-1	Seaside II	Pennlinks	Lopez	Penn A-4	Penn A-2	Penn A-1	Penn G-2	Southshore	Regent	Providence	SR 1020	Cato	Putter	Penn G-6	Penncross	Procup	Crenshaw	LSD (0.05)

¹Pure Seed Testing Research Farm, Hubbard, Oregon. Second Year Test.

Table 4. Varieties Significantly different from Penn A-1 for Listed Years 1995, 1996, 1997.

		Т		i	Τ	Τ	Т	Τ					Į	Ī	1		Ī		1		1
Total	Years	9	7	6	5	6	∞	∞	8	6	9	6	8	8	6	6	7	9	10	∞	
/horl	SS	97		97			97		97	26	26	97	26		97	76		97	76		Ī
wer W	Branches		96	96	96	96	96	8	96				96	96	96				96	96	5
# Lower Whorl	Bı			95								95							95	95	
							97			97			62	97	76				76		
Flag Leaf	Width	96		96										96							,
FI					95	95				E						95			95		
af		97		97	97	26	97	97			97	97	26		16	67	67	97	62	26	
Flag Leaf	Length		96																		30
Ē	-		95	95		95		95	,			95				95	95	95		95	
	ngth					26		97		62	62		26			16	97		26		
	Panicle Length	96	96	96	96	96	96		96	96	96	96		96	96	96	96	96	96	96	7 0
	Pani	95	95			95	95	95	95	95			95	95							
ive	ead																				
Vegetative	Base Spread																				1.0
ζę	Bas						95	95	95	95		95	95	95	95	95	56	95	95	95	
	ight					26		26	26	97		97			26	97	26			97	
	Plant Height	96	96	96		96			96	96	96	96		96	96	96	96	96	96	96	36
	Pla		95	95	95		95	95	95	95	95	95	95	95	95						
									50	0			e l			_			ę		1
	iety	Penn A-2	Penn A-4	Penn G-1	Penn G-2	Penn G-6	Seaside II	Seaside	Penncross	Penneagle	Pennlinks	er	Southshore	ent	SR 1020	Crenshaw	Jup.	ez	Providence		Potol Vocas
	Variety	Pen	Pen	Pem	Pen	Peni	Seas	Seas	Pen	Pen	Pem	Putter	Sou	Regent	SR	Cre	ProCup	Lopez	Prov	Cato	Total

Additional Description of Penn A-1

In close height of cut putting green management Penn A-1, along with codeveloped Penn A and G varieties, is unique in producing a very fine textured, dense high quality turf. Although closely related to Penn G-2, Pennlinks, and Lopez in spaced plant morphology; Penn A-1 differs in turfgrass qualities as follows:

Table 5	Turfgrass texture versus	Penn G-2: 7.8 vs 7.4 Pennlinks: 7.8 vs 6.7 Lopez: 7.8 vs 6.3
Table 6	Summer density versus	Penn G-2 7.7 vs 7.2 Pennlinks: 7.7 vs 6.2 Lopez: 7.7 vs 5.7
Table 7	Turfgrass quality versus	Penn G-2: 6.3 vs 6.1 Pennlinks: 6.3 vs 5.8 Lopez: 6.3 vs 5.6
Table 8	Fusarium patch versus	Penn G-2: 8.0 vs 6.8 Pennlinks: 8.0 vs 7.0 Lopez: 8.0 vs 7.5
Table 9	Dollarspot versus	Penn G-2: 7.6 vs 6.8 Pennlinks: 7.6 vs 7.5 Lopez: 7.6 vs 6.8
Table 10	Brownpatch versus	Penn G-2: 7.8 vs 7.5 Pennlinks: 7.8 vs 7.3 Lopez: 7.8 vs 7.4

LEAF TEXTURE RATINGS OF BENTGRASS CULTIVARS GROWN ON A GREEN 1994 DATA

7
FINE
9≃VERY
1-9;
RATINGS
TEXTURE
LEAF

	MEAN		7.8	7.8	7.7	7 2	- 1	4.	** <u>*</u>	7.4	2	. ~				· ·	Ξ,	:	6.7	6.7	6.7	6.7	6.5	4.9	6.3	6.3	6.3	, - -		0.0	٠. د.	5.6	2.5	2
	¥ F		3	89	8	×	1	-	B)	8.0	80	C	7 7	α		9 0	0 1	- 1	7.7	7.7	7.7	8.0	7.7	7.3	8.0	8.0	7.3	7.7	. a) (? ;) ·	?	0.7
	VA1	ł	3	: :	0	7	7	;	6.3	7.3	6.7	6.7	7.3	2.9	× ×	, ,) 0 1	2.	9.0	5.7	5.3	0		5.7	2.4	5.0	7		1 0		?	0.0
-	SC1	i	*	•	i,	2.7	7		0	0	5.7	5.7	4.7	10	7	7	, r	•) t	•	0.4	4.7	2.0	4.3	M M	4.0	5.0	M	7 7	~	, ,	- 10	;	1.2
1	RI1	•	*	01	0.	6.3	7		9 1	6.3	6.7	6.3	6,3	7.0	6.0	6.7	^			2 .	0.0	6.3	γ) ·	0.7	٧.0	2.7	5.7	6.0	2.7	· •	, ,	, ,	•	1.9
	PA1	` r			α. Ο.	7.7	7.7	7) r	:	7.3	7.7	8.0	7.0	7.3	7.7	7.0	7		: (٠ • •	٠. د) ·	4.	0. 1	 	0	M.	2.7	4	~		?	1.1
	oK1	c	Ł	- r	:	0.6	۳, ش	κ		۰ د د	8.7	7.3	8,3	8.0	8.7	8.3	7.7	7 7	. ^) ·	- r	- 1	2 1	? (0.7	7.7	7.0	7.3	6.7	7	9		6.0
	M02	α		o a	0.0	а. С	6.7	7.7	- c	9 0	o. 6	ο. Ο.	8 0	8 0.	7.7	8.0	7.7	8.0	7.7	. a	•	0 0	9 0	0 0) 0	י נים	÷:	O 1	7.7	7.3	8.0	7.3		7.0
	MN1	7.7	ŀ	7.7	. ,	?	7.7	7.3	2) c	? :	6.3	2.9	7.3	6.7	7.0	7.0	6.7				7	, v	י ה י	•) ·	? !	2.9	6.7	6.7	6.7		1.0
	K	8.7	, 20	7.7	. 0	- 1	8.	8.3	7.7	. ~	- r	? ;	 	 		7.7	8.0	7.0	7.0	7.0	2 2						o 6	? !	0.0	3.7	33	4.3		1.1
	11	8.7	9.0	0.6	2	. 0	> >	8	8.7	, M	2 0	? *		, i	· ·	α. Υ	8.7	8 0.	8,0	8.0	8	- C	C .	, c) a	• •	: 0) i	8.7	8.7	7.3		0.7
	A21	7.7	8.3	8	K	, ,	·	7.3	7.3	7.0	. ^	: ^	. v) ·	?!	٠. د د	6.7	2.7	2.4	6.0	5.7	, 15 15	6.0	N.	Y		. M			0.4	4.0	4.0		8.0
	NAME	A-1	A-4	SYN 92-1	9-9	CATO	2	Z-5	SYN 92-5	L-93	CRENSHAU	CVB 02-2	Sp 1020	BAD US 42402	20124 24 100	CONTRIBUTE	SOUTHSHORE	1S1-AP-89150	18TH GREEN	4-50	PENNL INKS	MSUEB	REGENT	PRO/CUP	LOPEZ	SYN-1-88	PENNCROSS	TRIES TAR	100000	I CADENZ	BAR AS 492	SEASIDE		LSD VALUE

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

DENSITY RATINGS 1-9; 9=MAXIMUM DENSITY 1/

				VULL	LICEUM DENSITY		•		
NAME	AZ1	GA1	M.	0K1	PA1	0E1	HA3	WAG	MEAN
PENN A-4 (A-4)	9.0	5.4	«	0	ν,	0	•	1	1
PENN A-1 CA-11	2) i	0 (⊃ ·	8.5	1.7	6.7
		?	ė	8.7	8.7	8.0	9.0	8.3	7.7
DENI O COLUMN OCCUPANTO	9	2	6.3	8.7	//	8.0	8.0	8.3	۲
		4.3	7.3	ω ~	8	8	7.3	8	7 2
PENN G-2 (G-2)	8.3	7	6.0	0.6	8	7.7	~	7	
CATO	7.7	4.3	6.3	8	7	α) C		
IMPERIAL (SYN 92-5)	7.7	7.7	2.7	7.3	7	, c	9 0		:
LOFT'S L-93 (L-93)	7.7	۷.0	6.3	8.7	7			, v	- 6
SR 1020	7.3	6.3	6.3	60	7.0	7.7	. ~	, c	9 0
BAR WS 42102	7.3	7 0	7.0	7.7	79	8			, ,
CRENSHAW	7.3	5.3	5	7.7	6.7	8	2.2		9 4
SYN 92-2	6.7	5.0	6.7	7.3	7 3	7.7	. ~		9 0
SOUTHSHORE	7.3	٤.3	5.7	7.7	9	, M			0 4
PROVIDENCE	7.3	1.7	5.3	7.0	7.7	7.7	7.0	, c	9 6
PENNL INKS	6.0	6.3	5,3	6.7	6.7	P.	2 4	, r	
ISI-AP-89150	6.7	0.4	5.7	7.0	7.3	0		ה	9 4
18TH GREEN	6.3	4.0	6.3	 	6.3	7.7	. 9	, w	4 -
d-50	5.7	4.0	7.4	6.7	6.7	7.7	6.7	;	· •
PRO/CUP	6.3	4.3	4.3	7.0	5.3	8	6.3	6.0	
TRUEL INE	5.7	4.3	5.3	7.0	2.7	7.7	2.0	, LO	, v
MSUEB	6.0	4.7	4.7	7.3	6.3	7.7	6.0	7.	, r.
LOPEZ	6.0	4.0	4.7	6.7	5.0	8.0	4.7	***	
PENNCROSS	6.3	5.0	3.7	5.7	5.0	7.7	4.7	6.7	
MARINER (SYN-1-88)	5.7	4.0	3.7	5,3	6.3	8.0	6.3	6	
REGENT	7.0	4.3	4.7	6.7	5.7	7.3	5	· ~	ı v
BAR AS 492	M.	6.0	5.0	2.3	3.7	7.0	~	. 7	
TENDENZ	4.0	4.7	•	2.7	7	2.7	9		, ,
SEASIDE	4.0	4.0	3.7	2.3	2.0	8	3.7	2.0	3.4
I SD VALUE	•	ć	•		,	•	ı		
ראף אינטב	<u>:</u>	·.	5.	2	0.	0.	2.0	4.9	0.5

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

MEAN TURFGRASS QUALITY RATINGS OF BENTGRASS CULTIVARS GROWN ON A GREEN AT TWENTY-SIX LOCATIONS IN THE U.S. 1994 DATA

TURFGRASS QUALITY RATINGS 1-9; 9=IDEAL TURF 1/

3 4 1 2	בראם האם	6.5	4.0	6.3	6.3	١	2				9	9	T.	. 0	. 60	т. Ф	, RO	5.7	5,7	5.7	9,	5.6	5.6	, ru	5,5	2.4	8,4	4.6	4.5	0
 S		۳. ا ا	5.	7.2	7.4	ŀ		. ^	, r.	1 4	, E	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	8	0	0.7	7.9	.3	6.7	5.5	7.0	6.7	9.9	9.9	6.1	9.9	6.3	8,5	5.5	5.6	. c
=						ı																							5.6	r.
. 2																													3.1	9 0
																													5.1	9
144	<u>.</u>	٥.٠		0.9	6.4																								3.8	5.0
Ž	<u> </u>	Ξ:	4.0		6.7	ŀ	0.9	2.9	0	8	5.5	6.9	6.5	7.2		7.2	6.0	5.7	6.5	6.1	4.9	6.5	6.5	7.0	6.1	6.5	6.1	5.2	6.3	1.2
ru.	; (7.	- :	4.3	5.6	2.0	5.0	7	5.1		7.7	7.7	4.4	4.6	4.7	4.3	3.0	5.0	4.0	4.0	3.4	3.6	3.8	3.5	3.4	3.8	3.1	2.5	2.8	8.0
11		- (0.0	5.8	5.9	6.5	8.5	6.1	6.2	5	5.4	5.7	6.1	5	5.4	5.6	5,2	5.2	6.9	5.2	5.6	6.4	4.8	5.6	4.8	3.8	5.0	4.8	6.0
		0 ^		δ.	7.3	<u> </u>	8.9	6.5	6.7	2.9	6.1	6.3	5.7	80	6.2	5.6	8.9	6.3	6.3	5.9	5.6	5.5	6.2	5.3	5.4	4.5	5.1	5.5	3.3	0.7
ZZ		0,0	† (٠	6.3	۶. ا	6.0	6.7	6.1	5.7	0.	5.0	in in	8	2.3	5.4	5.5	5.3	5.3	5.7	5.1	5.3	5.3	4.9		5. 5.	3.7	3.6	4.0	9.0
1.8		0.0	3 (7.0	7.2	6.5	7.7	9.9	6.8	5.9	ۍ ش	6.2	6.0	5.1	5.4	8.2	4.8	ۍ. ه	5.3	5.2	5.8	ς. Β	5.3	4.7	5.0	5.3	4.2	9.0	3.0	9.0
, H	,	- 0	, ,		5.2	7.9	6.7	7.9	ν. Ο	9	5	5.1	5. 89	4.9	6.0	6.2	5.9	5.3	5.6	7.2	5.6	6.0	5.8	6.2	6.3	5.0	5.0	2.7	3.6	0.4
, 5 <u>8</u>	1		, L	3	7.2	7:7	7.0	7.0	7.2	7.2	7.3	7.1	7.1	7.0	7.4	7.3	7.0	7.2	7.2	7.3	7.1	7.	7:	7.1	7.1	7.1	6.8	7.0	7.0	9.0
MO.		10		•	9.9	6.9	7.0	6.3	6.9	7.1	7.1	8.9	6.9	6.8	7.3	6.8	6.8	7.5	7.2	7.1	7.2	6.6	6.9	7.0	7.4	6.7	6.3	6.0	6.0	0.7
MN.	0			•	7.4	6.9	7.2	7.5	7.8	7.	7.6	6.5	7.3	7.1	6.8	7.0	7.4	9.9	7.2	7.5	8.8	6.7	7.0	6.8	7.0	6.8	6.5	6.3	4.9	8.0
Ξ	4	, v	, 0	0	5.2	5.8	5.8	ŗ,	5.8	5.5	5.2	υ, ες	N.	5,5	5.2	5.2	υ, M	5.2	5.2	5.3	5.2	4.8	7. 5.	5,2	4.3	4.7	4.5	3.5	4.0	9.0
MA 1			, ,	? !	7.3	7.0	7.7	7.5	6.5	6.1	9.9	7.9	6.0	5.9	5.8	6.2	5.7	5.6	9.9	6.3	6.1	2.7	6.1	5.2	N.	6.9	5.0	3.8	3.8	0.9
ΚY2	7 2	7		3	7.4	7.4	8.8	8.8	7.1	6.5	7.3	9.9	7.2	6.8	6.5	6.7	7.2	7.2	6.1	7.1	6.0	7.2	6.0	7.3	6.9	6.7	5.1	4.6	5. B	9.0
Κ¥	7 7	, c	,		Š.	7.7	8 0	7.7	7.8	7.1	7.	7.1	6.7	7.1	7.7	7.1	7.4	7.2	7.3	6.8	7.7	7.3	7.5	7.0	7.3	7.1	5.8	4.6	6.3	0.5
KS1	9	0	4		Ģ 0	5.	6.7	5.	5.4	9.9	7.0	5.6	6.5	6.1	6.5	6.5	6.5	6.0	ν. Ο		0.	9.9	2.7	5.0	4.9	5.0	4.3	5.6	6.9	1.4
11.2	7 1	4	4	;	?	9.	5.3	ນຸ	5.7	6.5	6.5	7.1	7.3	5.7	4.9	6.0	ις L	6.7	.0 0	4.7	6.3	 L	5.7	6.1	4.0		5. W	4.1	W.0	8.
=	ר. ה	, IV	0	, ,	ر د ز	2	5.1	4.8	6.7	6.7	 -	5.1	4.5	4.8	5.0	4.7	4.6	7.7	4.8	4.1	4.3	4.5	7.5	4.7	4.3	4.8	M	3.1	W. O.	9.0
IA1	2.9	9	4	:		٠ 8	9.9	6.3	5.4	6.2	ۍ ه	ۍ ش	2	9.6	ις 1,1	7	, M	5.4	'n,	5.2	ນ ນໍ	8.	4.7	N.	5.	4.3	3.4	3.6	3.5	0.0
GA2	2.1	0	7 0	, ,	7:5	5.5	2.0	2.5	2.1	6.	2.5	6.	2.3	2.4	2.5	1.8	7.	2.2	1.7	7.	3,2	1.7	.	2.7	2.5	2.0	4.4	5.	3.3	7.0
GA 1	0.4	6.3	6 7	, ,	ì	,	7	۴.1	3	4.5	3	9	7 7	4 3	4.3	~	٥ م	۲.1	3	3.7	M	۲,	o M	4.3	eo M	4.6	9	4.8	4.3	0.7
AZ1	7.2	6.6	6			3 1	7.3	6.2	6.2	6.0	6.1	5.7	2.7	υ. Ο	5.0	5,4	5. 0.	4.8	5.0	5.4	8.4	0.0	v. 0	4.9	4.8	4.8	2.5	2.4	3.7	0.7
NAME	A.54	* [-93	* PROVIDENCE	N-1			* CA10	9-9		* SOUTHSHORE	SYN 92-1	SYN 92-5	SYN 92-2	* SR 1020	* PENNLINKS	* REGENT	BAR WS 42102	MSUEB		* 18TH GREEN		* PRO/CUP				* SYN-1-88	* TENDENZ	BAR AS 492	* SEASIDE	LSD VALUE

COMMERCIALLY AVAILABLE IN THE USA IN 1995.

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). **/**

Table 8

FUSARIUM PATCH RATINGS OF BENTGRASS CULTIVARS GROWN ON A GREEN 1995 DATA

-		•
DISEASE	MEAN	88.88 60.00 6.77 7.77 7.77 7.77 7.77 7.77 7.7
9=N0	WA3	88.37 7.77 88.37 7.75 7.75 7.75 7.75 7.75 7.75 7.75 7
RATINGS 1-9;	WA1	88.0 6.0 7.7 7.7 7.7 7.7 7.7 7.7 7.7 7.7 7.7 7
FUSARIUM PATCH RATI	NAME	CENTURY (SYN 92-1) PROVIDENCE PENN A-1 (A-1) TENDENZ BAR WS 42102 PENN A-4 (A-4) LOFI'S L-93 (L-93) REGENT SOUTHSHORE BAR AS 492 CATO IMPERIAL (SYN 92-5) LOPEZ SR 1020 SRN 92-2 18TH GREN ISI-AP-89150 PRO/CUP CRENSHAW GG-P PENN G-6 (G-6) PENN G-6 (G-6) PENN G-6 (G-6) PENN G-6 (G-6) PENNLINKS TRUELINE MSUEB

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). -

1.6 1.1

LSD VALUE

DOLLAR SPOT RATINGS OF BENTGRASS CULTIVARS GROWN ON A GREEN 1995 DATA

	DOLL	DOLLAR SPOT RATINGS 1-9; 9=NO DISEASE	RATINGS	1-9;	P=NO DIS		1.		
NAME	KS1	MN	MO1	N.51	о <u>к</u>	PA1	1X	VA1	MEAN
TENDENZ	7		•	(1				
BAR AS 492	7.0	٠,	9 6	8 5	8.3	0	8.0	8.0	8.1
10FT 1 - 92 (1 - 02)	- 6	? !	2.0	8.7	8.3	0	œ 0.	6.7	~
? :		7.0	8	7.7	8.0	7.7	M		9 6
FENN A- ((A-1)	7.3	8.0 0.0	8.0	7 1	<u>د</u>	7 7	, ,	: '	۰.۱
PENNL INKS	\. 8	٥	ķ	ŀ	-	1	3	ç.	9.7
MSUEB	7.7	,			ر د د	0 1	0.7	7.3	5.
CATO			0 1	0.	8.7	7.3	۲.	6.3	7.4
SEASIDE			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	6.3	8.3	7.7	8.3	7.0	7 2
DENNOOCE) · ;	0.	7.7	6.7	8.3	8.0	M.		7.
TENACROSS SPECIAL SPECIAL SPEC	8.0	0.9	7.7	7.7	8.7	7.7	2.9		
FRUVIDENCE 10: 10 00/10	7.3	7.7	7.0	6.7	7.7	2.9		. ^	• •
151-AP-89150	7.7	7.3	7.0	6.7	~	· ·		? (y (
d-50	7.3	6.0	7.7	4				٠ • •	0.7
REGENT	7.3	6.3	4	•	, ,	9 6	- 1	0	7.0
BAR WS 42102	٧	7	, ,		2.6) ·	۳. د	9.0	7.0
LOPEZ	7		• •	ر د د	יי יי	O	7.0	6.0	7.0
PENN G-2 (G-2)		? ;		٠,	œ.	6.7	7.3	5.7	8.9
		: ;	0°9	6.3		7.3	7.3	4.7	8,9
CONTRETIONS			6.7	6,3	8.3	6.7	7.0	6.0	×
SUCH SHOKE	0.0	κ, i	6.7	2.7	8.0	6.7	0.0	7.0	2 6
	5.0	0.7	5.3	9.0	8.0	6.3	7.7	7	¥ ¥
PENN A-4 (A-4)	2	6.3	7.3	5.7	8.3	6.3	2		, 4
PRU/LUP	0 /	5.7	6.3	5.7	8.0	6.3	7.0	7.7	
MAKINEK (STN-1-88)	6.3	7.3	0	5.7	8.0	10°	7.7	, c	יי ני טיי
IMPERIAL (SYN 92-5)	5.0	7.3	6.3	0.4	7.7	2	~ ~	, ,	? 4
SR 1020	۲.	6.3	6.3	4.7	~	۲ ۲	, ,) i
SYN 92-2	5	4.7	6.3	~	. ~) h		?!	0 1
CENTURY (SYN 92-1)	7.2) Y	,	? !	י י	9 1	5.5	5.3
18TH GREEN	'n	, ,	? <	? ;		. ·	~	بر بر	4.7
CDENCHAL		- !	.	7	6.5	2.7	7.3	6.3	4.6
	÷	.	5.0	2.3	6.7	3.7	6.7	3.3	4.5
LSD VALUE	1.3	2.0	1.6	1.2	-	,	u	•	
				!	:		?	†	٠ <u>.</u>

1/ TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05).

Table 10

BROWN PATCH RATINGS OF BENTGRASS CULTIVARS GROWN ON A GREEN 1994 DATA

BROWN PATCH RATINGS 1-9; 9=NO DISEASE 1/

MEAN	880.7 000.8 8 8 8 7 7 7 8 8 8 8 7 7 8 8 8 8 8 8 8	1.0
sc1	0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	ж Т.
PA1	88 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	
0K1	88.00 7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.7.	1.2
N.J	88.0 7.7.7 88.8 8.7.7 8.8 8.7.7 8.8 7.7.7 8.8 8.0 8.0 8.0 8.0 7.7.7 8.0 8.0 8.0 8.0 8.0 8.0 7.7.7 8.0 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 7.7.7 8.0 8.0 8.0 7.0 7.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8.0 8	1.3
NAME	MSUEB 18TH GREEN G-6 A-1 CRENSHAU TRUELINE L-93 SOUTHSHORE PRO/CUP CATO G-2 PRO/CUP CATO G-2 PRO/IDENCE REGENT LOPEZ SYN 92-2 SYN 92-2 SYN 92-2 SYN 92-2 SYN 92-1 ISI-AP-89150 PENUCROSS SEASIDE BAR WS 42102 TENDENZ BAR AS 492	LSD VALUE

TO DETERMINE STATISTICAL DIFFERENCES AMONG ENTRIES, SUBTRACT ONE ENTRY'S MEAN FROM ANOTHER ENTRY'S MEAN. STATISTICAL DIFFERENCES OCCUR WHEN THIS VALUE IS LARGER THAN THE CORRESPONDING LSD VALUE (LSD 0.05). =

REPRODUCE LOCALLY. Include form number and date on all reproductions. U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE SCIENCE AND TECHNOLOGY DIVISION - PLANT VARIETY PROTECTION OFFICE		ede in accordance with the Privacy A sperwork Reduction Act (PRA) of 1991
EXHIBIT E STATEMENT OF THE BASIS OF OWNERSHIP	Application is consisted in auto-	to determine if a plant variety prote
1. NAME OF APPLICANT(S)	2. TEMPORARY DESIGNATION OR EXPERIMENTAL NUMBER	3. VARIETY NAME
Pennsylvania Agricultural Experiment Station	A-1	Penn A-1 Creeping Bentgras
A ADDRESS (Sec.)		orosping benegras
4. ADDRESS (Street and No., or R.F.D. No., City, State, and ZIP Code, and Country) Charles R. Krueger Bruce McPheron Associate Dean	5. TELEPHONE (include area code)	6. FAX (include area code)
Associate Dean	814-865-5410	814-863-7905
0217 Agricultural Admin. Bldg.	7. PVPO NUMBER	1014 003 7303
University Park, PA 16802	97000	. 4.2
8. Does the applicant own all rights to the variety? Mark an "X" in appropriate is	<u> </u>	145
	orock. It no, piease explain.	X YES NO
9. Is the applicant (individual or company) a U.S. national or U.S. based company	?	
If no, give name of country		X YES NO
O. Is the applicant the original breeder? If no, please answer the following:		
 a. If original rights to variety were owned by individual(s): ls (are) the original breeder(s) a U.S. national(s)? If no, give name of contractions 	ountry	YES X NO
		_
b. If original rights to variety were owned by a company: Is the original breeder(s) U.S. based company? If no, give name of cour		x YES NO
 Additional explantion on ownership (If needed, use reverse for extra space): J. M. Duich, Professor Turfgrass Science is breed Employee of applicant with rights assigned to applicant. 	der. plicant.	
EASE NOTE:		
nt variety protection can be afforded only to owners (not licensees) who meet one	e of the following criteria:	
If the rights to the variety are owned by the original breeder, that person must be of a country which affords similar protection to nationals of the U.S. for the same		OV member country, or national
If the rights to the variety are owned by the company which employed the original nationals of a UPOV member country, or owned by nationals of a country which a genus and species.		be U.S. based, owned by nals of the U.S. for the same
If the applicant is an owner who is not the original breeder, both the original breed	der and the applicant must meet	one of the above criteria
original breeder may be the individual or company who directed final breeding	g. See Section 41(a)(2) of the	Plant Variety Protection Act for
reporting burden for this collection of information is estimated to average 10 minutes per response, including the data needed, and completing and reviewing the collection of information. Send comments regarders stions for reducing this burden, to Department of Agriculture, Clearance Officer, OIRM, AG Box 7630, James 20055 and form number in your letter.	ing the time for reviewing instructions, searing this burden estimate or any other aspect is L. Whitten Buildian Washington D.C.	thing existing data sources, gathering and of this collection of information, including
		va.v. veikil reprying, refer to UMB No.
the PRA of 1995, no persons are required to respond to a collection of information unless it displays a valid O		